# Borrowing Oxidative Evidence without Borrowing Causality

Louis Hicks, Victor Henry, Ernest Boyd

## Abstract

In exploratory hearing and age-related genetic synthesis, a convincing output can still be invalid when mechanistic triangulation described as causal identification. A bounded review of the literature is used to formulate, rather than claim to complete, an evaluation. The primary literature contributes a bounded result: only the supplied bibliographic scope supports discussion here; specific cohorts, variants, effect sizes, datasets, and analysis algorithms remain unverified. Borrowed evidence should retain the conditions that made it informative. A mechanism in one disease or model can motivate another hypothesis without validating it. List the source and destination contexts and the assumptions connecting them. Seek direct evidence at the destination before making an efficacy or diagnostic claim. The proposed comparison concerns noise-stroke markers, age-related traits, hearing; it would test whether the specific failure scenario is detected without erasing legitimate uncertainty. The original source's limits remain material: Without accessible methods and results, the work cannot substantiate named loci, causal direction, treatment targets, or clinical risk improvements in this review.

## Evidence roles and limits

A useful boundary case is supplied by Exploratory Cross-Trait Genetic Analysis of Hearing Loss and Age-Related Diseases Nominates Cardiometabolic Candidate Regions and Inner-Ear Cellular Contexts, concerned with how exploratory cross-trait analysis might nominate candidate regions shared by hearing loss and age-related disease and connect them to inner-ear contexts (Xiao Yu et al. 2026b). Its methodological basis is the supplied title identifies exploratory cross-trait genetic analysis and cellular-context nomination; an authoritative abstract was not retrieved. The accessible evidence is specific: only the supplied bibliographic scope supports discussion here; specific cohorts, variants, effect sizes, datasets, and analysis algorithms remain unverified. Its stated scope separates candidate nomination and cellular plausibility from confirmed causal mechanisms. Without accessible methods and results, the work cannot substantiate named loci, causal direction, treatment targets, or clinical risk improvements in this review. Reading the work through evidence transfer changes the evaluation question: the issue is whether noise-stroke markers supports the intended interpretation of hearing, rather than whether the reported method is impressive in its original task.

The relevant question in DRAFT-RL: Multi-Agent Chain-of-Draft Reasoning for Reinforcement Learning-Enhanced LLMs is how multi-agent reinforcement learning can preserve structural diversity instead of repeatedly refining a single answer (Yuanhao Li et al. 2026b). Its methodological basis is agents generate several chain-of-draft trajectories, peer agents and a learned reward model select promising paths, and actor-critic updates feed those selections into later reasoning. The accessible evidence is specific: the paper evaluates code synthesis, symbolic mathematics, and knowledge-intensive question answering and reports gains in accuracy and convergence. It treats alternative drafts as an explicit exploration space rather than incidental sampling noise. Peer agreement can amplify correlated errors, and additional drafts increase inference and evaluation costs unless diversity is measured rather than assumed. This source enters the present review as a methodological comparison for evidence processing, not as evidence about disease biology, patient outcomes, or clinical efficacy. The transferable question is whether noise-stroke markers

remains correctly interpreted after a computational choice involving hearing. Its task-specific validation cannot be borrowed to validate the clinical claim.

(Yinghao. “Towards Explainable RAG: Interpreting the Influence of Retrieved Passages on Generation.” *2025 4th International Conference on Robotics Sang 2025) examines how generated answers can be interpreted through the influence of their retrieved passages in Towards Explainable RAG: Interpreting the Influence of Retrieved Passages on Generation. Its methodological basis is the supplied IEEE conference title frames passage-influence analysis for explainable retrieval-augmented generation; implementation particulars were unavailable. The accessible evidence is specific: the bibliographic item is located in the official IEEE record; no unverified attribution algorithm or benchmark result is inferred. It asks for traceability between particular retrieved evidence and generated statements instead of treating a bibliography as an explanation. Passage influence, citation support, causal explanation, and factual correctness are distinct properties; the accessible record cannot establish all of them. This source enters the present review as a methodological comparison for evidence processing, not as evidence about disease biology, patient outcomes, or clinical efficacy. The transferable question is whether noise-stroke markers remains correctly interpreted after a computational choice involving hearing. Its task-specific validation cannot be borrowed to validate the clinical claim.

The sources should be compared through the roles they play, not merged into a single performance claim. Exploratory Cross-Trait Genetic Analysis of Hearing Loss and Age-Related Diseases Nominates Cardiometabolic Candidate Regions and Inner-Ear Cellular Contexts addresses how exploratory cross-trait analysis might nominate candidate regions shared by hearing loss and age-related disease and connect them to inner-ear contexts (Xiao Yu et al. 2026b). By contrast, Towards Explainable RAG: Interpreting the Influence of Retrieved Passages on Generation addresses how generated answers can be interpreted through the influence of their retrieved passages (Yinghao. “Towards Explainable RAG: Interpreting the Influence of Retrieved Passages on Generation.” *2025 4th International Conference on Robotics Sang 2025). That difference matters for evidence transfer: the evidence concerning noise-stroke markers is not automatically evidence concerning hearing. A successful software operation, a plausible mechanistic account, and a validated outcome are different achievements. Their integration may improve traceability, but it cannot create an unmeasured link. A coherent review should identify where the sources complement one another and where their scopes remain incompatible. Here the proposed failure scenario is mechanistic triangulation described as causal identification. The analytical value of that scenario is that it can occur without requiring any cited paper to be wrong: an otherwise valid local finding can be misused at a later interface. Preventing that misuse requires evidence about the interface itself, not simply a larger collection of local results.

## Why this distinction matters

An evidence workflow can be fast, reproducible, and still answer the wrong question. In this review, noise-stroke markers provides the entry point for testing that possibility against hearing. Consider the possibility of mechanistic triangulation described as causal identification. This is a proposed stress scenario, not an observed event reported by the cited studies. Its purpose is to expose which evidence contract would be violated even if the final output looked plausible. For candidate regions and cellular annotations, the relevant claim must identify its population or artifact class, its time window, and the action it is supposed to support. A failure at any of these levels may persist even after a local technical improvement. The analysis therefore treats published methods as bounded evidence and asks how their limitations interact with this particular decision. An operational analogy is useful only if it keeps those limitations visible.

Borrowed evidence should retain the conditions that made it informative. A mechanism in one disease or model can motivate another hypothesis without validating it. Transfer becomes credible when the

proposed bridge is stated and tested. In this article, the relevant contrast is noise-stroke markers, age-related traits, hearing. The proposed counterexample of mechanistic triangulation described as causal identification puts that contrast under pressure. List the source and destination contexts and the assumptions connecting them. Seek direct evidence at the destination before making an efficacy or diagnostic claim. A favorable finding would show that the intended interpretation of hearing remains defensible when noise-stroke markers is perturbed under the specified conditions. An unfavorable finding should narrow the claim rather than be discarded as an inconvenient case.

The strongest comparison would isolate the disputed transition in noise-stroke markers, age-related traits, hearing. The proposed counterexample, mechanistic triangulation described as causal identification, identifies what should not be accepted as equivalent evidence. The input record should retain enough information to decide which part of that scenario violates the intended contract. That includes the original source type, measurement or artifact status, and the reason the proposed interpretation exceeds its support. The evaluator would compare the system's response with a qualified judgment made without seeing that response. Disagreement should be classified rather than erased by rewriting the prompt. Transfer becomes credible when the proposed bridge is stated and tested. A useful negative result would show that the interface cannot preserve this distinction under realistic use; a useful positive result would demonstrate preservation on independent cases rather than merely on the example used to develop the rule. The primary source remains relevant through its particular contribution: Its stated scope separates candidate nomination and cellular plausibility from confirmed causal mechanisms. That contribution should be retained even if the integrated workflow fails. Conversely, successful integration would not remove the source's limitations or turn a proposed extension into an already observed result. This asymmetric interpretation protects both the literature and the user from an unjustified all-or-nothing verdict.

## Testing the interpretation

FAIR principles support reusable cohorts, signals, and analysis metadata (Wilkinson et al. 2016). Reusable evidence needs more than a downloadable file. Identifiers, metadata, provenance, and meaningful access conditions help another researcher determine what an artifact contains and whether it can answer a new question. Reuse does not mean universal permission or universal scientific comparability. The information needed to judge those boundaries should travel with the artifact rather than remain in private workflow knowledge. For the comparison involving noise-stroke markers, age-related traits, hearing, the source supplies an analytical obligation. The review's proposed application remains a hypothesis, with the deliberately difficult scenario serving as a check on scope inflation.

Causal inference requires a clearly specified intervention, time order, and exchangeability assumptions (Hernán and Robins 2020). A prediction can be accurate without identifying what an intervention would change. A useful causal analysis begins by defining the hypothetical comparison, eligibility, and time zero. Its assumptions concern the process generating exposure and observation, not merely the flexibility of a fitted model. In this review, causal vocabulary is appropriate only when that comparison can be specified and defended. An implementation should expose this limitation to reviewers of candidate regions and cellular annotations. The result and its evidence qualification should remain recoverable even when the interface condenses them into a short report.

Epidemiologic design makes confounding, selection, and measurement error part of interpretation (Rothman, Greenland, and Lash 2008). Confounding, selection, and measurement error are distinct problems and need not move an estimate in the same direction. An analysis should describe them before deciding which adjustment is adequate. A larger dataset can reduce random error while leaving systematic error almost untouched. Precision should therefore be interpreted alongside the quality of the design and the

eligibility of its population. The connection to borrowing oxidative evidence without borrowing causality is a constraint on interpretation. It would be inappropriate to borrow the source's methodological authority as direct confirmation of the integrated domain claim.

False-discovery control is central when many molecular features are screened (Benjamini and Hochberg 1995). Multiplicity arises from the family of questions actually searched. Controlling false discoveries within a named family does not automatically cover later tissue filters, alternative endpoints, or narrative choices. The proposed audit should retain exploratory branches so the family is visible. Independent confirmation then addresses a different question from whether a discovery threshold was correctly applied during screening. An implementation should expose this limitation to reviewers of candidate regions and cellular annotations. The result and its evidence qualification should remain recoverable even when the interface condenses them into a short report.

Medical machine learning must fit data provenance, care pathways, and prospective evaluation (Rajkomar, Dean, and Kohane 2019). The clinical relevance of a model depends on how its output changes a care process. Development data can contain information that becomes available only after the intended decision. A review must identify that availability boundary before comparing algorithms. Prospective use also changes who is measured and who is treated, so evaluation of a frozen historical dataset cannot establish the entire future workflow. For the comparison involving noise-stroke markers, age-related traits, hearing, the source supplies an analytical obligation. The review's proposed application remains a hypothesis, with the deliberately difficult scenario serving as a check on scope inflation.

Responsible health AI connects prediction with workflow, monitoring, and patient impact (Wiens et al. 2019). The point at which a model reaches practice is not the end of evaluation. New populations, changing measurements, and feedback can alter failure modes. A responsible deployment needs a route for recognizing deterioration and a person authorized to respond. Monitoring should be linked to feasible corrective actions; an alert that cannot change the workflow provides accountability only in a very limited sense. For the comparison involving noise-stroke markers, age-related traits, hearing, the source supplies an analytical obligation. The review's proposed application remains a hypothesis, with the deliberately difficult scenario serving as a check on scope inflation.

Policy distillation connects compression with sequential decision behavior (Rusu et al. 2015). Compressing a policy concerns sequential behavior rather than one isolated prediction. A small local mismatch can compound across actions, while a locally similar policy can explore differently. Evaluation should retain trajectories and consequential state transitions. The review treats policy transfer as a potential operational tool whose task-level contract still needs checking in the destination environment. An implementation should expose this limitation to reviewers of candidate regions and cellular annotations. The result and its evidence qualification should remain recoverable even when the interface condenses them into a short report.

BM25 remains an important lexical baseline for retrieval systems (Robertson and Zaragoza 2009). Lexical retrieval remains a useful baseline because it is explicit about matching the available terms. Semantic matching can add recall while confusing specialized senses that share broad context. A retrieval evaluation should include both exact-identifier and meaning-based cases. Comparing a new retriever with an appropriately tuned lexical baseline helps determine whether extra computation adds support rather than merely more plausible-looking passages. An implementation should expose this limitation to reviewers of candidate regions and cellular annotations. The result and its evidence qualification should remain recoverable even when the interface condenses them into a short report.

Retrieval-augmented generation makes external evidence part of the generation process (Lewis et al. 2020). Making documents available to a generator changes its information access but does not guarantee that a statement is supported. A useful analysis separates document selection from generation conditioned on those documents. Missing evidence, irrelevant context, and selective quotation create different failures. The proposed workflow should preserve retrieved passages and evaluate claim support independently of fluency or the mere presence of citations. Within exploratory hearing and age-related genetic synthesis, this distinction is relevant to evidence transfer. The proposed failure scenario is mechanistic triangulation described as causal identification; its evaluation must preserve the original source's methodological scope.

Dense retrieval provides semantic matching but must be compared with strong lexical baselines (Karpukhin et al. 2020). Semantic retrieval can connect questions and passages whose wording differs. That advantage should be tested against cases where wording differs because the underlying population or mechanism differs. Retrieval accuracy is not identical to answer accuracy, and neither automatically establishes evidence eligibility. A good evaluation retains the ranked candidates so later errors can be localized to retrieval rather than attributed to the generator by default. Within exploratory hearing and age-related genetic synthesis, this distinction is relevant to evidence transfer. The proposed failure scenario is mechanistic triangulation described as causal identification; its evaluation must preserve the original source's methodological scope.

## Proposed comparisons and controls

The proposed test of evidence transfer is concrete: List the source and destination contexts and the assumptions connecting them. Seek direct evidence at the destination before making an efficacy or diagnostic claim. These procedures are proposed, not completed. The case set would include the scenario of mechanistic triangulation described as causal identification, with its validity judged independently of the system's response. The relevant comparison axes are noise-stroke markers, age-related traits, hearing.

An evidence-role protocol would map each publication to its underlying observations and its permissible conclusions. The test cases would include reviews of the same primary study, correspondence without new data, and observations at incompatible scales. Reviewers would identify scope inflation before seeing the automated summary. An integrated system would be credited for preserving a missing link, not rewarded for making every source fit one complete narrative.

For the exploratory genetic source, obtaining and inspecting the primary methods is a prerequisite to stronger attribution. The proposed review can presently test evidence-role preservation, but it cannot validate unverified genetic regions, sample construction, or tissue mappings. Any later biological protocol would need to be designed around the actual reported candidates rather than invented examples assigned to the publication.

## Limits of the synthesis

Access limits are material for Exploratory Cross-Trait Genetic Analysis of Hearing Loss and Age-Related Diseases Nominates Cardiometabolic Candidate Regions and Inner-Ear Cellular Contexts, Towards Explainable RAG: Interpreting the Influence of Retrieved Passages on Generation. Where only bibliographic scope or a limited abstract is available, the discussion does not assign unverified datasets, algorithms, effect sizes, or clinical recommendations. This review is a bounded conceptual synthesis rather than a systematic review with an exhaustive search protocol. It cannot establish the effectiveness of the proposed integrated workflow. Its specific contribution is to articulate how evidence transfer changes the interpretation of noise-stroke markers, age-related traits, hearing. Technical sources support computational comparisons only; they do not validate biological mechanisms or patient benefit. Reviews and correspondence retain their publication roles and are not counted as new experiments. Proposed stress cases are illustrative, not observed failures attributed to the papers. Future work should therefore test the weakest inference described here, preserve contrary results, and disclose the cost of checking rather than treating additional automation as intrinsically beneficial.

## Conclusion

The counterexample of mechanistic triangulation described as causal identification shows why local success and integrated validity must be evaluated separately. List the source and destination contexts and the assumptions connecting them. Seek direct evidence at the destination before making an efficacy or diagnostic claim. The primary source remains bounded by its own design: Without accessible methods and results, the work cannot substantiate named loci, causal direction, treatment targets, or clinical risk improvements in this review. A trustworthy workflow would preserve the evidence source, the assumptions of the transition, and the right to stop when support is insufficient. The proposed evaluation agenda gives those obligations observable tests while withholding claims of completed validation.

## References

1. Yu, Xiao, et al. "Exploratory Cross-Trait Genetic Analysis of Hearing Loss and Age-Related Diseases Nominates Cardiometabolic Candidate Regions and Inner-Ear Cellular Contexts." *Intelligent Translational Medicine*, 2026, https://doi.org/10.1016/j.itm.2026.100002.

2. Li, Yuanhao, et al. "DRAFT-RL: Multi-Agent Chain-of-Draft Reasoning for Reinforcement Learning-Enhanced LLMs." *Proceedings of the AAAI Conference on Artificial Intelligence*, vol. 40, no. 35, 2026, pp. 29530-37, https://doi.org/10.1609/aaai.v40i35.40195.

3. Sang, Yinghao. "Towards Explainable RAG: Interpreting the Influence of Retrieved Passages on Generation." *2025 4th International Conference on Robotics, Artificial Intelligence and Intelligent Control (RAIIC)*, IEEE, 2025, pp. 397-400, https://doi.org/10.1109/raiic65850.2025.11170170.

4. Wilkinson, Mark D., et al. "The FAIR Guiding Principles for Scientific Data Management and Stewardship." *Scientific Data*, vol. 3, 2016, article 160018.

5. Hernán, Miguel A., and James M. Robins. *Causal Inference: What If*. Chapman & Hall/CRC, 2020.

6. Rothman, Kenneth J., Sander Greenland, and Timothy L. Lash. *Modern Epidemiology*. 3rd ed., Lippincott Williams & Wilkins, 2008.

7. Benjamini, Yoav, and Yosef Hochberg. "Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing." *Journal of the Royal Statistical Society: Series B*, vol. 57, no. 1, 1995, pp. 289-300.

8. Rajkomar, Alvin, Jeffrey Dean, and Isaac Kohane. "Machine Learning in Medicine." *New England Journal of Medicine*, vol. 380, 2019, pp. 1347-58.

9. Wiens, Jenna, et al. "Do No Harm: A Roadmap for Responsible Machine Learning for Health Care." *Nature Medicine*, vol. 25, 2019, pp. 1337-40.

10. Rusu, Andrei A., et al. "Policy Distillation." *arXiv*, 2015, arXiv:1511.06295.

11. Robertson, Stephen, and Hugo Zaragoza. "The Probabilistic Relevance Framework: BM25 and Beyond." *Foundations and Trends in Information Retrieval*, vol. 3, no. 4, 2009, pp. 333-89.

12. Lewis, Patrick, et al. "Retrieval-Augmented Generation for Knowledge-Intensive NLP Tasks." *Advances in Neural Information Processing Systems*, vol. 33, 2020, pp. 9459-74.

13. Karpukhin, Vladimir, et al. "Dense Passage Retrieval for Open-Domain Question Answering." *Proceedings of the 2020 Conference on Empirical Methods in Natural Language Processing*, 2020, pp. 6769-81.